

XV. Follow-up schedule for nonseminoma stage I, surveillance

Name:	Civic registration number:			
Orchiectomy, date:	_Side:	right / left	LVI: yes/no	
Date definitive staging:	_			
Recurrence date:	_fill out	fill out new form and switch FU		

TUMOR MARKERS SHOULDBE CHECKED EVERY 2 MONTHS YEAR 1 FOR PATIENTS WITH LVI+ MANAGED BY SURVEILLANCE

Control type **B**: Patient contact, AFP, β -hCG, LDH, S-creatinine, **MRI of the retroperitoneum**/

(abdominopelvic CT).

Control type **C**: Like B with addition of clinical examination, testosterone, SHBG, LH, FSH. Control type **TM**: Tumour markers, AFP, β-hCG and LDH (*List the patient for a telephone*

rumour markers, AFF, p-ned and LDH (List the putient joi

appointment).

Scrotal ultrasound when clinically indicated. Metabolic screening (lipids, fasting glucose, HbA1c), and blood pressure at 1- and 5-year visit. Inform Swedish patients at 1- and 5- year visit that a quality of life questionnaire will be sent out from RCC Syd, Sweden.

Months from end of treatment:

	TM	В	TM	<u>c</u>	Follow-up year 1
0	3	6	9	12	
	TM	В	TM	В	Follow-up year 2
12 15	18	21	24		
		TM		<u>C</u>	Follow-up year 3
24		30		36	
		TM		В	Follow-up year 4
36		42		48	
		TM		<u>C</u>	Follow-up year 5
48		54		60	

Postpubertal teratoma: No TM check-up month 3, 9, 15, 21

Patient care plan to be given to the patient at termination of follow-up